

foodsafe Allergy Test



NEW PROFILE INCLUDES

- **95 FOODS TESTED**
Includes most common food allergens.
- **NO BLOOD DRAW**
This lab test requires only a small amount of blood taken from a finger-prick and is very easy to do.
- **CLINICALLY RELEVANT**
IgG food allergy testing for delayed reactions. About 60% of all undiagnosed symptoms are due to food allergies.*
- **ROTATION DIET PLAN**
Included with every food allergy report.
- **FAST. EASY. AFFORDABLE.**

FINDING HIDDEN FOOD ALLERGIES JUST GOT A LOT EASIER FOR YOUR PATIENTS

*About 60% of all undiagnosed symptoms are due to food allergies according to James Breneman, MD, former Chairman of the Food Allergy Committee of the American College of Allergy (now the American College of Allergy and Immunology), in his 1978 book, Basics of Food Allergy.

This is a test for delayed hypersensitivity reactions. Delayed (IgG) responses may cause delayed symptoms, such as joint or muscle pain, chronic headaches, fatigue, eczema, and psoriasis. Since IgG reactions often occur hours or days after particular foods are consumed, correlating symptoms can be difficult. The reactions can be subtle or severe, and may lead to chronic (long-term) symptoms and chronic degenerative conditions.



FOODSAFE™
Allergy Test

Meridian Valley
Laboratory

6839 Fort Dent Way, Ste. 206 Tukwila, WA 98188 206.209.4200

WWW.MERIDIANVALLEYLAB.COM